

(51) International Patent Classification 5 : A61K 7/00, 7/06, 7/08		A1	(11) International Publication Number: WO 90/10429 (43) International Publication Date: 20 September 1990 (20.09.90)
(21) International Application Number: PCT/US90/00802 (22) International Filing Date: 12 February 1990 (12.02.90) (30) Priority data: 323,480 13 March 1989 (13.03.89) US (71) Applicant: S.C. JOHNSON & SON, INC. [US/US]; Patent Section, M.S. 077, 1525 Howe Street, Racine, WI 53403-5011 (US). (72) Inventor: UICK, Heide, J. ; 139 Greshell Lane, Racine, WI 53406 (US). (74) Agents: RAKOCZY, Richard, E. et al.; Patent Section M.S. 077, S.C. Johnson & Son, Inc., 1525 Howe Street, Racine, WI 53403-5011 (US).		(81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), IT (European patent), LU (European patent), NL (European patent), SE (European patent). Published <i>With international search report.</i>	
(54) Title: SPARKLING PEARLESCENT PERSONAL CARE COMPOSITIONS			
(57) Abstract			
<p>This invention relates to cationic oil-in-water emulsion compositions which are especially useful as hair conditioning compositions which have a sparkling pearlescent appearance due to the presence of irregular platelet-like crystals suspended in the composition. The compositions comprise a cationic water-in-oil emulsion of from (a) about 0.1 % to 5 %, preferably from 0.1 to 0.8 %, of a quaternary ammonium compound such as stearyl dimethyl benzyl ammonium chloride; (b) about 1-10 %, preferably 1-2 %, of a fatty acid having 12-16 carbon atoms such as myristic acid and a fatty monoalcohol having from 12 to 18 carbon atoms such as cetyl alcohol in a 70:30 to 95:5, preferably 75:25 to 85:15, weight ratio of acid to alcohol; (c) about 0.5-2 % of a compatible thickening agent such as hydroxyethyl cellulose; optionally, up to about 5 % of additional cationic, nonionic and amphoteric surfactants; and the balance comprises water wherein the pH of the emulsion is from about 2-5, preferably from 2-4, and the viscosity is preferably from about 1 to 5 Pascal.seconds (1,000 to 5,000 centipoise) at 25°C.</p>			
BEST AVAILABLE COPY			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	HU	Hungary	NO	Norway
BJ	Benin	IT	Italy	RO	Romania
BR	Brazil	JP	Japan	SD	Sudan
CA	Canada	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LI	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark				

SPARKLING PEARLESCENT PERSONAL CARE COMPOSITIONS

5

TECHNICAL FIELD

This invention relates to personal care compositions such as hair conditioning compositions which are cationic water-in-oil emulsions which have a sparkling pearlescent appearance due to the presence of irregular platelet-like crystals.

BACKGROUND ART

Pearlescing agents are sometimes included in personal care compositions such as hair conditioners and shampoos to make them more attractive to the consumer. These conventional agents include ethylene glycol monostearate, ethylene glycol distearate, guanine, bismuth oxychloride on mica and the like. Suspended fish scales (guanine) can give a sparkling effect to such compositions, but are dependent on natural sources. U.S. Pat. No. 4,654,207 to Preston teaches apearlescent shampoo where the pearlescing agent is a fatty acid ester like myristyl myristate or cetyl myristate which is added to the shampoo base from a substantially anhydrous solubilizing agent such as a surfactant. The pearlescent effect is observed after about 5 days waiting time which is unlike the direct use of myristyl myristate taught in Preston's earlier U.S. Pat. No. 4,438,096. Columns 1-2 of the '207 Patent further discuss the state of the art relative to pearlescing agents in cosmetic compositions.

FIGURE 1 of this Specification is a photomicrograph at 400X magnification of a commercially available shampoo product sold by the Procter and Gamble Company under the name "PERT(R) Shampoo" which contains glycol distearate, i.e., ethylene glycol distearate. The pearlescing agent produces a uniform pearlescence to produce a shiny uniform sheen.

U.S. Pat. No. 3,320,133 to Suga et al. teaches a pearlescent face lotion. A phenol compound and a monohydric alcohol are said to be required to get floating pearlescent crystals of fatty acids in the lotion. Example 1 has 5 parts of stearic acid, 2 parts of linoleic/linolenic acid and 0.5

parts of decyl alcohol (93.3:7.7 ratio of fatty acid to fatty alcohol). Example 2 has no decyl or other fatty alcohol, but is said to be pearly. Example 3 has ethoxylated/propoxylated cetyl alcohol, but no other fatty alcohols. None of the 5 examples contain quaternary ammonium salts, but all do contain significant amounts of alkanolamines to obtain a composition of weak alkalinity. This differs from my invention as will be explained further below.

U.S. Pat. No. 3,590,122 to Hutcheson et al. does not 10 indicate that the conditioning shampoos it teaches are pearlescent. The product is composed of an organic synthetic detergent plus a saturated fatty acid like isostearic acid that is neutralized at least in part. Adjuvants such as fatty alcohols (e.g., cetyl alcohol or stearyl alcohol) can 15 be present in an amount of up to 5% by weight, but the only reason for inclusion given is to "impart desired qualities".

A number of patents describe using fatty alcohols and fatty acids in hair conditioners and shampoos, some of which further contain fatty alkyl quaternary ammonium compounds:

20	<u>U.S. Pat. No.</u>	<u>Inventor</u>	<u>Comment</u>
	2,968,628	Reed	Examples 3 and 11
	3,149,042	Habicht et al.	Example 4
	3,502,769	Fukuhara	Example 2
	4,374,125	Newell	Example 3, also has quaternary
25	4,551,330	Wagman et al.	Formula B has quaternary, cetyl alcohol and stearic acid (6/94 acid/alcohol)
	4,608,392	Jacquet et al.	Examples 8, 15 and 16 have stearic acid/cetyl alcohol, highest is 75/25
30	4,614,200	Hsiung et al.	Example 3
	3,651,931	Hsiung	Example 3 has 95.7/3.3 oleic acid/lauryl alcohol
	4,777,039	Lang et al.	Pearlescent conditioner with quaternary plus fatty alcohol, no fatty acid
35	4,357,141	Grollier et al.	Hair dyes with polymeric quaternary plus fatty acid (oleic) and fatty alcohol

<u>U.S. Pat. No.</u>	<u>Inventor</u>	<u>Comment</u>
		(octyldodecyl) at 2/1 or lower acid/alcohol ratios
4,201,766	Grollier et al.	Polymeric quaternary, optionally has quaternary compound, Formulas 2g, 3c & 3g; col. 9 teaches fatty acids and fatty alcohols

Further representing the state of the art are U.S. Pat. Nos. 3,155,591 (Hilfer); 4,007,261 (Sorrentino et al.); 4,278,657 (Tezuka et al.); 4,345,080 (Bolich); 4,421,740 (Burton); 4,478,853 (Chaussee); 4,493,824 (Abe); and 4,701,322 (Dixon et al.).

15 SUMMARY DISCLOSURE OF THE INVENTION

After studying the effects of including fatty acid in hair conditioning compositions which are cationic water-in-oil emulsions, I have found that the use of a relatively high ratio of certain fatty acids to certain fatty alcohols in the presence of certain fatty alkyl quaternary ammonium compounds unexpectedly results in a composition which has a sparkling pearlescent appearance which results in an attractive product. The sparkling nature of the compositions is due to the presence of irregular platelet-like crystals which tend to turn as the composition is poured or shaken. These crystals are illustrated in FIGURE 2 of this Specification.

These compositions do not require the use of inorganic platelet-containing materials such as mica or fish scale type materials such as guanine.

30 These and other advantages are provided by a composition which comprises a cationic water-in-oil emulsion of from (a) about 0.1% to 5%, preferably from 0.1 to 0.8%, of a quaternary ammonium compound such as stearyl dimethyl benzyl ammonium chloride; (b) about 1-10%, preferably 1-2%, of a fatty acid having 12-16 carbon atoms such as myristic acid and a fatty monoalcohol having from 12 to 18 carbon atoms such as cetyl alcohol in a 70:30 to 95:5, preferably 75:25 to 85:15, weight ratio of acid to alcohol; (c) about 0.5-2% of a compatible thickening agent such as hydroxyethyl cellulose;

optionally, up to about 5% of additional cationic, nonionic and amphoteric surfactants; and the balance water wherein the pH of the emulsion is from about 2-5, preferably from 2-4, the viscosity is preferably from about 1 to 5 pascal-seconds 5 ("Pa.s") (1,000 to 5,000 centipoise) at 25°C. wherein the composition contains irregular platelet-like crystals which impart a sparkling appearance to the composition.

BRIEF DESCRIPTION OF THE DRAWINGS

10 In the accompanying Drawings:


FIGURE 1 is a photomicrograph using cross-polarized light at 400X magnification of a commercial pearlized shampoo composition.

15 FIGURE 2 is a photomicrograph at 400X magnification of the composition of the present invention showing the irregular platelet-like crystals responsible for the sparkling appearance of this composition.


DETAILED DESCRIPTION OF THE PRESENT INVENTION

20 This invention relates to a composition having a sparkling pearlescent appearance comprising a cationic oil-in-water emulsion of

a) from about 0.1% to 5% based on the total weight of the emulsion of at least one cationic surfactant selected 25 from the group consisting of quaternary ammonium compounds of the formula



30 $[R^1R^2N^+(R^5)_2]A^-$; and



- where R^1 is an alkyl group having an average of 8 to 23 carbon atoms, R^2 is an alkyl group of 1 to 4 carbon atoms, R^3 is R^1 or R^2 , R^4 is R^2 or a benzyl group, R^5 is $(CH_2CH_2O)_nH$ where n is an integer of from 1 to 50, and A is an anion wherein the first formula does not include compounds where R^2 , R^3 and R^4 each have from 1 to 4 carbon atoms;
- 5 b) from about 1 to 10% by weight based on the total weight of the emulsion of
- 10 i) a fatty acid having an average of from 12 to 16 carbon atoms per molecule and
- ii) a fatty monoalcohol having an average of from 12 to 18 carbon atoms per molecule in a 70:30 to 95:5 by weight ratio of (i) to (ii);
- 15 c) from 0.5 to 2.0% by weight based on the total weight of the emulsion of a compatible thickening agent; and
- d) the balance comprises water,
- 20 wherein the pH of the emulsion is from about 2 to 5 and characterized in that the composition contains irregular platelet-like crystals which provide the sparkling appearance as a result of the presence of the surfactant (a) and the ingredients (b).
- 25 The present compositions are characterized by the presence of irregular platelet-like crystals with shiny surfaces which provide a sparkling appearance. The irregularity causes the crystals to tumble in the emulsion and the platelet-like character causes the shiny surfaces to
- 30 catch light and sparkle. These crystals do not plate out and form a uniform shine as is typical of many compositions containing pearlescing agents. FIGURE 1 is a color photomicrograph at 400X magnification using cross-polarized lighting of a product formerly sold by Procter & Gamble
- 35 Company as original "PERT(R) Shampoo for Normal & Dry Hair" which has a label ingredient statement as follows: Water, Sodium Laureth Sulfate, SD Alcohol 40, Cocamide MEA, Myristic Acid, Glycol Distearate, Fragrance, Disodium Phosphate, EDTA, Sodium Hydroxide, Imidazolidinyl Urea, Methylparaben,

Propylparaben, D&C Green No. 8, and FD&C Blue No. 1. This anionic shampoo formulation appears to employ Glycol Distearate as the pearlizing agent and FIGURE 1 shows that no significant amount of large crystals are evident in the composition. The pearlescent agent apparently plates out and forms a uniform shiny, pearlescent surface.

FIGURE 2 is a color photomicrograph at 400X magnification using cross-polarized lighting of a hair conditioning composition of the present invention which shows the presence of large crystals in various planes of rotation. Some surfaces of the crystals appear to be reflecting light towards the camera (i.e., towards the consumer's eye) while others are not. This alternating reflectance provides the sparkling appearance characteristic of these compositions. The irregularity of the crystals is also evident and this apparently prevents the crystals from plating out and forming a completely uniform pearlescence.

The compositions of the present invention are cationic water-in-oil emulsions which are especially useful as hair conditioning compositions. As such, they require the presence of at least one cationic surfactant in an amount of from 0.1% to 5% based on the total weight of the emulsion. More preferably, these surfactants comprise from about 0.1 to 0.8% of the emulsion composition and, most preferably from about 0.15 to 0.35% of the composition. I have found that certain such surfactants in the form of fatty alkyl quaternary ammonium compounds are necessary to cause the development of the crystals which provide the sparkling appearance noted above.

The required fatty alkyl quaternary ammonium surfactants are selected from the formulas noted as (a) above and are well known in the art, many of which are commercially available. Examples of surfactants of the formula $[R^1R^2N^+R^3R^4]A^-$ are those such as stearyl dimethyl benzyl ammonium chloride and lauryl dimethyl benzyl ammonium chloride, distearyl dimethyl ammonium chloride, dicoco dimethyl ammonium chloride and ditallow dimethyl ammonium chloride, and methyl tri(octyl-decyl) ammonium chloride which is sold under the tradename ADOGEN 464 by Sherex Chemical

Company, Inc. of Dublin, Ohio (hereinafter "Sherex"). This formula excludes formulas where R^2 , R^3 and R^4 each have from 1 to 4 carbon atoms. In these and the other formulas, A is an anion of the type conventionally used with quaternary ammonium compounds such as halides such as chloride and bromide, and alkyl sulfates such as methyl sulfate and ethyl sulfate, among others.

Examples of surfactants of the formula $[R^1R^2N^+(R^5)_2]A^-$ are ethyl bis (polyethoxy ethanol) alkyl ammonium ethyl sulfate where the R^1 alkyl group has from 8 to 18 carbon atoms and the size of the alkyl group and the value of n in the R^5 groups are such that the average molecular weight of the compound is about 1110 (VARSTAT (R) 66 from Sherex), methyl bis (polyethoxy ethanol) alkyl ammonium chloride where the R^1 alkyl group has from 8 to 18 carbon atoms and the size of the alkyl group and the value of n in the R^5 groups are such that the average molecular weight of the compound is about 910 (VARIQUAT (R) K-1215 from Sherex), and methyl bis (2-hydroxyethyl) coco ammonium chloride (VARIQUAT 638 from Sherex).

Examples of the last formula listed under (a) above are those where R^1 has from 12 to 18 carbon atoms such as methyl-1-tallow amido ethyl-2-tallow imidazolinium-methyl sulfate (VARISOFT (R) 475 from Sherex) and methyl-1-oleyl amido ethyl-2-oleyl imidazolinium-methyl sulfate (VARISOFT (R) 3690 from Sherex).

Most preferred are quaternary ammonium compounds of the formula $[R^1R^2N^+R^3R^4]A^-$ and, of these, preferably R^1 has an average of 18 carbon atoms, R^2 and R^3 are each alkyl groups having 1 carbon atom, R^4 is a benzyl group, and A is a chloride anion, i.e., stearyl dimethyl benzyl ammonium chloride.

Other surfactants which are compatible with the other ingredients present in the composition can be included such as conventional cationic, nonionic and amphoteric surfactants. The term "compatible" as used herein indicates that such ingredients do not cause the composition to become unstable or prevent the formation of the desired crystals to produce the sparkling effect. Preferably, no more than about

- 5% of such surfactants are included where a hair conditioning composition is desired and more preferably, no more than about 2% of additional compatible surfactants are present. These surfactants can be employed to improve the emulsion formed as well as to act as hair conditioners or anti-static agents, or to serve other beneficial purposes. Preferred additional surfactants are cationic surfactants such as fatty alkyl quaternary ammonium compounds and fatty amines which can be used for their hair conditioning properties.
- 10 Examples of such additional surfactants are quaternary ammonium compounds such as those of the formula $[R^1N^+(R^2)_3]A^-$ where R^1 , R^2 and A are as defined above and preferably, R^2 is a methyl group. These are exemplified by palmityl trimethyl ammonium chloride and coco trimethyl ammonium chloride.
- 15 Other cationic surfactants can be fatty alkyl amines such as stearyl dimethyl amine. Examples of nonionic and amphoteric surfactants can be polyethoxylated fatty alcohols, fatty ethoxylated esters of sorbitol and fatty amine oxides.

- Besides the required cationic surfactants of (a) above,
- 20 two other required ingredients are a total of from about 1 to 10% by weight based on the total composition, preferably from about 1 to 2%, of (i) a fatty acid having an average of from 12 to 16 carbon atoms per molecule such as lauric acid, myristic acid and palmitic acid and (ii) a fatty monoalcohol
- 25 having an average of from 12 to 18 carbon atoms per molecule such as lauryl alcohol, myristyl alcohol, cetyl alcohol and stearyl alcohol. Since fatty acids and alcohol are often derived from natural sources, there is generally a distribution of isomers present in the compositions and an
- 30 average carbon chain length is often used.

- To obtain the sparkling crystal appearance, the ratio of fatty acid to fatty alcohol must be from 70:30 to 95:5 and, more preferably, from 75:25 to 85:15 of acid to alcohol. Outside of these ratios, the crystals do not attain the
- 35 structure necessary to cause a rather noticeable sparkling appearance due to the presence of irregular platelet-like crystals. Although stearic acid was observed to form small crystals which reflected some light, it was not found to produce compositions having this desirable appearance.

To keep the crystals in suspension and obtain the desired viscosity, preferably between about 1 and 5 Pa.s (1,000 and 5,000 centipoise) at 25°C., from about 0.5 to 2% of a compatible conventional thickening agent of the type commonly used in cosmetic formulations is included in the compositions of the present invention. As used herein, "viscosity" refers to viscosity as measured on a Brookfield LV Viscosimeter using a #3 or #4 spindle, 30 r.p.m. for 30 seconds at 25°C. Examples of such thickeners are hydroxyethyl cellulose of the type sold under the tradename NATROSOL by Aqualon Company of Wilmington, Delaware as well as hydroxypropyl cellulose and other nonionic or cationic water soluble and water dispersible thickening agents. The exact composition of the thickening agent or protective colloid is unimportant as long as it is cosmetically acceptable and compatible with the composition.

Water comprises the balance of the composition. Although tap water can be used if it contains relatively few ions, deionized water is preferred.

Adjustment of the pH of these compositions is done in a conventional manner using cosmetically acceptable organic or inorganic acids with organic acids such as citric acid being preferred. The pH of the cationic emulsions of the present invention is in the range of 2 to 5 and is more preferably in the range of 2 to 4.

As further additional ingredients, conventional additives such as perfumes, fragrances, dyes, proteins, solubilizing agents such as propylene glycol, and the like can be added to the composition in relatively small amounts provided that they are compatible with the compositions.

INDUSTRIAL APPLICABILITY

These compositions can be prepared in a manner similar to that employed for hair conditioning emulsions. U.S. Pat. No. 4,421,740 to Burton gives one procedure for accomplishing the same and refers to equipment that can be employed. The object is to form an emulsion composition which is then aged for about 5 days to a month or more at room temperature

(about 20-25°C) to permit formation of the desired crystals in the composition.

Thus, the compositions of the present invention can be made by forming an intermediate composition which is generally about 25-30% of the total composition by mixing or dispersing together the cationic surfactant (a), any additional surfactant which may be used, the fatty acid, the fatty alcohol, pH adjusting agent, and sufficient water to permit the addition of 25-30 parts of this intermediate to 70-75 parts of additional ingredients to result in a total of 100 parts of the compositions of the present invention. The intermediate composition is heated to about 65-72°C. to assist in blending the ingredients together. After the ingredients are well-mixed, the composition is then cooled to about 43°C. At this temperature, any perfumes to be used are added and dispersed into the intermediate composition. This cooled mixture is then homogenized at 3.45 - 55.2 MegaPascal ("MPa") (500-8,000 p.s.i.), preferably at about 27.6 MPa (4,000 p.s.i.). Subsequent to the homogenization step, the intermediate is cooled back to about 25-35°C.

In a separate vessel, the remaining water and any dyes and preservative to be added is heated to about 40-45°C. with agitation and the thickening agent is added into the vessel with stirring. Stirring is continued until the thickening agent is hydrated and well-dispersed within the water. Any additional ingredients are then added with agitation to the vessel containing the hydrated thickening agent followed by 25-30 parts of the intermediate per 70-75 parts of the mixture in the vessel to make a total of 100 parts. Mixing is continued until a homogeneous composition is obtained. That composition is then aged at room temperature until such time as the composition has a sparkling pearlescent appearance due to the presence of irregular platelet-like crystals. Aging can take from about 5 days to one or more months. Aging can be conducted simply by packaging the composition in individual containers and storing the packaged compositions for a period of time.

The resulting compositions are especially suitable for use as ready to use hair conditioners or creme rinses which

are applied directly to the hair of the user subsequent to shampooing.

The following examples are provided to show various aspects of the present invention without departing from the scope and spirit of the invention. Unless otherwise indicated, all parts and percentages used in the Examples are by weight.

Example 1

This Example shows the preparation of a composition of the present invention which employs the preferred 80:20 ratio of myristic acid to cetyl alcohol and possessed a sparkling pearlescent appearance due to the presence of irregular platelet-like crystals. This is the composition shown in FIGURE 2.

The intermediate had the following composition:

Intermediate 1

Stearyl dimethyl benzyl ammonium chloride ¹	1.34
Cetyl alcohol	0.938
20 Myristic acid ²	3.752
Cetyl trimethyl ammonium chloride ³	1.34
Stearyl dimethyl amine ⁴	0.70
Citric acid	0.5075
Deionized water	90.5475
25 Fragrance	<u>0.875</u>
Total	100.00%

1. AMMONYX (R) 4002 from Stepan Company of Northfield, IL, 94% active.
- 30 2. EMERY 655 from Emery Division of Quantum Chemical Corp. of Cincinnati, OH.
3. VARIQUAT (R) E-228 from Sherex, 25% active.
4. ADOGEN (R) 342-D from Sherex.

The final composition had the following composition:

35

Composition 1

Deionized water	70.05
Preservative ¹	0.03
Hydroxyethyl cellulose ²	0.75

Collagen protein hydrolyzates	0.10
Propylene glycol	0.50
Intermediate 1	28.57
Total	100.00%

5 -----

1. KATHON (R) CG solution from Rohm & Haas Co. of Philadelphia, PA which is 1.5% of 5-chloro-2-methyl-4-isothiazolin-3-one and 2-methyl-4-isothiazolin-3-one.
2. NATROSOL (R) 250HHR from Aqualon.
- 10 3. LEXEIN (R) X250 from Inolex Chemicals Co. of Philadelphia, PA.

Intermediate 1 was prepared by mixing the ingredients in the order listed in a vessel with agitation followed by heating to 65 - 71°C. and holding at that temperature for about 15 minutes. The mixture was cooled to 43°C., the fragrance was added and agitation was continued for another 10 minutes. Intermediate 1 was then homogenized in a Matin-Gaulin Homogenizer at 27.6 MPa (4,000 p.s.i.) and held at 43°C. until it was used to prepare Composition 1.

- 20 Composition 1 was prepared by adding the preservative to the water in a vessel separate from the one containing Intermediate 1. The vessel contents were heated to 40-43°C. with agitation and then the hydroxyethyl cellulose thickening agent was sifted into the agitating contents of the vessel.
- 25 Agitation was continued until the thickening agent was fully hydrated and dispersed into the water. The protein and propylene glycol were then added while agitation was continued, followed by the addition of Intermediate 1. The composition was stirred an additional 15-30 minutes until it
- 30 was homogeneous. Upon cooling to 25°C., the composition had a viscosity of 2.48 Pa.s (2,480 centipoise) (#3 spindle, 30 r.p.m. for 30 seconds) and a pH of 2.65. After 51 days of storage at room temperature, Composition 1 had a pearlescent, sparkling appearance throughout the composition. A
- 35 subsequent batch using this formulation was noted to exhibit a sparkling appearance within about 20 days after preparation of the batch.

Example 2

In this Example, the formulation of Example 1 was modified by changing the ratio of myristic acid/cetyl alcohol to determine where compositions containing a sparkling pearlescent appearance would be obtained. Except for the changes in ratios, the formulations and procedures used to make them remained the same as in Example 1. In the listing on the following page, only the amount of myristic acid and cetyl alcohol used in the Intermediate is listed.

10 Intermediate

Myristic Cetyl

	<u>Acid</u>	<u>Alc.</u>	<u>Ratio</u>	<u>Viscosity</u>	<u>pH</u>	<u>Results</u>
	0.00	4.69	0:100	5,960* (5.96)	3.0	No platelets noted
15	1.876	2.814	40:60	3,120 (3.12)	2.60	No platelets noted
20	2.814	1.876	60:40	2,276 (2.276)	2.65	No platelets noted
	3.283	1.407	70:30	3,052 (3.052)	2.65	Small crystal platelets observed after 15 weeks; no change after 10 months
25	3.517	1.173	75:25	2,792 (2.792)	2.65	Crystal platelets noted after 5.3 months.
30	3.986	0.704	85:15	2,388 (2.388)	2.65	Small crystal platelets seen after 7 days obvious large platelets seen after 13 days
35	4.221	0.469	90:10	2,824 (2.824)	2.55	Crystal platelets noted after 6 days
	4.455	0.235	95:5	2,272 (2.272)	2.65	Crystal platelets noted after 6 days, definite platelets: 9 months
40	4.69	0.00	100:0	3,560 (3.56)	2.60	Crystals noted after 6 days, no change after 10 months

45 * - Brookfield LV, spindle #4, 30 r.p.m. at 30 seconds, remainder are #3 spindle, 30 r.p.m. at 30 seconds, 25°C. First Value is in centipoise, second value in parentheses is in Pa.s.

As shown above, no platelets were noted until a 70:30
50 ratio of myristic acid to cetyl alcohol was used and even

then, the platelets were not as large or as prominent as in the compositions with 75:25 through 95:5 ratios. About the same results obtained for a 70:30 ratio were observed at a 100:0 ratio of myristic acid (i.e., no cetyl alcohol present) although some crystal platelets were noted in the 70:30 ratio composition. In these evaluations, "platelets" or "crystal platelets" is intended to mean a significant amount of irregular platelet-like crystals which gives a noticeable sparkling effect. In this and other Examples, "Crystals" indicates that slight sparkling was seen due to many small crystals, but was not nearly as noticeable upon the casual observation that a consumer would make as opposed to compositions containing "crystal platelets", the latter being the most desirable.

Example 3

In this example, other types of fatty acids and fatty alcohols were substituted into the formulation used in Example 1 and prepared in the same manner. The types and amounts of these ingredients are reported below as well as the results of testing the compositions.

<u>Intermediate</u>						
	<u>Fatty Acid</u>	<u>Fatty Alc.</u>	<u>Ratio</u>	<u>Viscosity</u>	<u>pH</u>	<u>Results</u>
25	C ₁₀ : 3.752	C ₁₆ : 0.938	80:20	720 (0.72)	2.40	No platelets noted.*
30	C ₁₂ : 3.752	C ₁₆ : 0.938	80:20	1,540 (1.54)	2.50	A few platelets observed after 1 month, crystal platelets after 5 months.
35						
	3.986	0.704	85:15	1,100 (1.10)	2.55	A few platelets observed after 1 month, crystal platelets after 4 months.
40						
	C ₁₆ : 3.752	C ₁₆ : 0.938	80:20	3,360 (3.36)	2.35	Small platelets observed after 2 months, crystal platelets after 8 months.
45						

	<u>Intermediate</u>					
	<u>Fatty</u>	<u>Fatty</u>		<u>Viscosity</u>	<u>pH</u>	<u>Results</u>
	<u>Acid</u>	<u>Alc.</u>	<u>Ratio</u>			
5	3.986	0.704	85:15	3,504 (3.504)	2.55	Small platelets observed after 1 month, crystal platelets after 4.5 months.
10	4.69	0.00	100:0	3,420 (3.42)	2.55	No platelets observed.
15	C ₁₈ : C ₁₆ : 3.752	0.938	80:20	3,784 (3.784)	2.35	Crystals observed after 2 months, no change after 9 months.
20	3.986	0.704	85:15	3,768 (3.768)	2.55	Very small platelets/ crystals observed after 2.5 wks., no change after 4.5 months.
25	4.69	0.00	100:0	2,560 (2.56)	2.50	Very small platelets/ crystals observed after 2 months, no change after 6.75 months.
30						
35	C ₁₄ : C ₁₂ : 3.752	0.938	80:20	3,696 (3.696)	2.50	Crystal platelets noted after 7 days.
40	C ₁₄ : C ₁₄ : 3.752	0.938	80:20	3,552 (3.552)	2.50	Crystal platelets observed after 2.5 months.
45	4.221	0.469	90:10	5,320** (5.32)	2.35	Crystal platelets observed after 3.5 months.
50	C ₁₄ : C ₁₈ : 3.752	0.938	80:20	2,076 (2.076)	2.55	A few platelets noted after 7 days, crystal platelets observed after 4 months.

55 * - Sample separated into 2 phases after 3 weeks.

** - Brookfield LV, spindle #4, 30 r.p.m. at 30 seconds, remainder are #3 spindle, 30 r.p.m. at 30 seconds, 25°C.

First value is in centipoise, second value in parentheses is in Pa.s.

Fatty acids:

- C₁₀: Capric acid - EMERY 659 from Emery.
 5 C₁₂: Lauric acid - EMERY 652 from Emery.
 C₁₄: Palmitic Acid - EMERSOL 143 from Emery.
 C₁₈: Stearic Acid - EMERSOL 153 from Emery.

Fatty Alcohols:

- C₁₂: Lauryl alcohol - EPAL 12 from Ethyl Corp. of
 10 Baton Rouge, LA.
 C₁₄: Myristyl alcohol - Lanette 14 from Henkel.
 C₁₈: Stearyl Alcohol - ADOL (R) 62 NF from Sherex.

Of the fatty acids tested, capric acid and stearic
 15 acid did not produce compositions containing the type of
 crystal platelets needed for the present invention. Use of
 palmitic acid without any fatty alcohol did not produce
 compositions containing the type of crystal platelets needed
 for the present invention. The three fatty alcohols tested
 20 all produced compositions containing the type of crystal
 platelets needed for the present invention in the ratios
 tested.

Examples 4-12

25 These Examples show the results of using different
 quaternary ammonium compounds or amounts thereof to produce
 compositions of the present invention. These compositions
 were prepared using the following formulations according to
 the procedure given in Example 1:

30 Intermediates

<u>Example No.:</u>	<u>4</u>	<u>5</u>	<u>6</u>	<u>7</u>
AROSURF (R) TA-1001	1.696	--	--	--
35 ADOGEN (R) 4642	--	1.876	--	--
VARISOFT (R) 4753	--	--	2.098	--
VARSTAT (R) 664	--	--	--	2.214
Cetyl alcohol	0.938	0.938	0.938	0.938
Myristic acid	3.752	3.752	3.752	3.752
40 ADOGEN (R) 342-D	0.700	0.700	0.700	0.700
Citric acid	0.5075	0.5075	0.5075	0.5075
Deionized water	91.5315	91.3515	91.1295	91.0135
Fragrance	0.875	0.875	0.875	0.875
Total	100%	100%	100%	100%

-
1. Distearyl dimethyl ammonium chloride (94%) from Sherex.
 2. Methyl tri(C8-C10) ammonium ethyl sulfate (85%) from Sherex.
 - 5 3. Methyl-1-tallow amido ethyl-2-tallow imidazolinium methyl sulfate (76%) from Sherex.
 4. Ethyl bis (polyethoxy ethanol) alkyl ammonium ethyl sulfate (72%) from Sherex.

10

Intermediates

Example No.:	8	9	10	11
VARIQUAT K-12151	1.678	--	--	--
15 AMMONYX (R) 4002	--	1.696	--	--
VARISOFT (R) 1102	--	--	2.126	--
VARIQUAT E-228	--	--	--	6.378
Cetyl alcohol	0.938	0.938	0.938	0.938
Myristic acid	3.752	3.752	3.752	3.752
20 ADOGEN (R) 342-D	0.700	0.700	0.700	0.700
Citric acid	0.5075	0.5075	0.5075	0.5075
Deionized water	91.5495	91.5315	91.1015	86.8495
Fragrance	0.875	0.875	0.875	0.875
Total	100%	100%	100%	100%

25

-
1. Methyl bis (polyethoxy ethanol) alkyl ammonium chloride (95%) from Sherex.
 2. Methyl bis (hydrogenated tallowamidoethyl) 2-hydroxyethyl ammonium methyl sulfate (75%) from Sherex.

30

To prepare the final compositions, the following formulation was used as in Example 1 where the intermediate employed corresponded to those described above: 70.05% deionized water, 0.03% KATHON (R) CG, 0.75% NATROSOL (R) 250HHR, 0.10% LEXEIN (R) X-250, 0.50% propylene glycol and 28.57% Intermediate (described above).

Example 12 was the same as Example 1, except the perfume, protein and propylene glycol components were not included. Thus, the Intermediate used to prepare Example 12 was as follows: 1.34% AMMONYX (R) 4002, 0.938% cetyl alcohol, 3.752% myristic acid, 1.34% VARIQUAT (R) E-228, 0.70% ADOGEN 342-D, 0.5075% citric acid and 91.4225% deionized water. This intermediate was homogenized at 20.7 MPa (3,000 p.s.i.) due to an inability to obtain 27.6 MPa (4,000 p.s.i.) pressure.

The final composition for Example 12 had the formula: 70.65% deionized water, 0.03% KATHON (R) CG, 0.75% NATROSOL (R) 250HHR, and 28.57% Intermediate (described above).

Example 13 was similar to Example 12, but one half of the amount of the AMMONYX (R) 4002, VARIQUAT (R) E-228 and ADOGEN (R) 342D were used. Thus, the Intermediate used to prepare Example 13 was as follows: 0.67% AMMONYX (R) 4002, 0.938% cetyl alcohol, 3.752% myristic acid, 0.67% VARIQUAT (R) E-228, 0.35% ADOGEN 342-D, 0.5075% citric acid and 93.1125% deionized water. The final composition for Example 13 had the formula: 70.65% deionized water, 0.03% KATHON (R) CG, 0.75% NATROSOL (R) 250HHR, and 28.57% Intermediate (described above).

Example 14 was similar to Example 12, but the AMMONYX (R) 4002 and VARIQUAT (R) E-228 were eliminated from the formulation and only the ADOGEN (R) 342D was used. Thus, the Intermediate used to prepare Example 14 was as follows: 0.938% cetyl alcohol, 3.752% myristic acid, 0.70% ADOGEN 342-D, 0.5075% citric acid and 94.1025% deionized water. The final composition for Example 14 had the formula: 70.65% deionized water, 0.03% KATHON (R) CG, 0.75% NATROSOL (R) 250HHR, and 28.57% Intermediate (described above).

Example 15 was similar to Example 12, but the AMMONYX (R) 4002 was eliminated from the formulation. Thus, the Intermediate used to prepare Example 13 was as follows: 0.938% cetyl alcohol, 3.752% myristic acid, 1.34% VARIQUAT (R) E-228, 0.70% ADOGEN 342-D, 0.5075% citric acid and 92.7625% deionized water. The final composition for Example 13 had the formula: 70.65% deionized water, 0.03% KATHON (R) CG, 0.75% NATROSOL (R) 250HHR, and 28.57% Intermediate (described above).

The finished compositions were evaluated and the results were as follows:

Ex. No.	Viscosity	pH	Results
35 4	2,388 (2.388)	2.50	Fine crystal platelets noted after 1.4 months.
5	2,072 (2.072)	2.45	Crystal platelets noted after 1.4 months.
40 6	1,088 (1.088)	2.65	Fine crystal platelets noted after 5 weeks.

	<u>Ex. No.</u>	<u>Viscosity</u>	<u>pH</u>	<u>Results</u>
	7	928 (0.928)	2.70	Very fine crystal platelets noted after 5 weeks.
5				
	8	1,484 (1.484)	2.80	Very slight crystal platelet formation noted after 1 month.
10	9	2,320 (2.320)	2.55	Fine crystal platelets noted after 2.5 weeks, crystal platelets noted after 6 wks.
15	10	3,124 (3.124)	2.75	No platelets formed after 1 month.
	11	2,824 (2.824)	2.35	No platelets formed after 2 months.
20	12	1,984 (1.984)	2.45	Crystal platelets noted after 3 weeks.
	13	2,724 (2.724)	2.40	Crystal platelets noted after 1 month.
25				
	14	2,368 (2.368)	2.65	No platelets formed after 2 months.
30	15	1,944 (1.944)	2.45	No platelets formed after 2 months.




Viscosity - Brookfield LV, #3 spindle, 30 r.p.m. at 30 seconds, 25°C. First value is in centipoise; second value in parentheses is in Pa.s.

35 Examples 4-7, 12 and 13 formed or were appearing to form crystal platelets at the time of the last observation. At the last observation, Example 8 appeared to need more time to form crystal platelets. Examples 10, 11, 14 and 15 were comparative examples which did not form crystal platelets.

What I claim is:

1. A composition having a sparkling pearlescent appearance comprising a cationic oil-in-water emulsion of

- a) from about 0.1% to 5% based on the total weight of the emulsion of at least one cationic surfactant selected from the group consisting of quaternary ammonium compounds of the formula



where R^1 is an alkyl group having an average of 8 to 23 carbon atoms, R^2 is an alkyl group of 1 to 4 carbon atoms, R^3 is R^1 or R^2 , R^4 is R^2 or a benzyl group, R^5 is $(CH_2CH_2O)_nH$ where n is an integer of from 1 to 50, and A is an anion wherein the first formula does not include compounds where R^2 , R^3 and R^4 each have from 1 to 4 carbon atoms;

- b) from about 1 to 10% by weight based on the total weight of the emulsion of
- i) a fatty acid having an average of from 12 to 16 carbon atoms per molecule and
 - ii) a fatty monoalcohol having an average of from 12 to 18 carbon atoms per molecule in a 70:30 to 95:5 by weight ratio of (i) to (ii);
- c) from 0.5 to 2.0% by weight based on the total weight of the emulsion of a compatible thickening agent; and
- d) the balance comprises water,

wherein the pH of the emulsion is from about 2 to 5 and characterized in that the composition contains irregular platelet-like crystals which provide the sparkling appearance as a result of the presence of the surfactant (a) and the
5 ingredients (b).

2. The composition of Claim 1 which further includes up to 5% of at least one additional compatible surfactant selected from the group consisting of cationic surfactants, nonionic surfactants and amphoteric surfactants.

10 3. The composition of Claim 1 wherein (b) is present in an amount of from about 1 - 2%, the ratio of (i) to (ii) is from 75:25 to 85:15, the fatty acid has an average of about 14 carbon atoms and the fatty alcohol has an average of about 16 carbon atoms per molecule.

15 4. The composition of Claim 1 wherein said (a) is $[R^1R^2N^+R^3R^4]A^-$ and is present in an amount of from about 0.1 to 0.8%.

5. The composition of Claim 4 wherein the composition contains up to about 2% by weight of at least one compatible
20 cationic surfactant, has a pH in the range of from about 2 to 4 and has a viscosity at 25°C. of from 1 to 5 Pascal.seconds (1,000 to 5,000 centipoise).

6. The composition of Claim 5 wherein (b) is present in an amount of from about 1 - 2% and the ratio of (i) to
25 (ii) is from 75:25 to 85:15.

7. The composition of Claim 6 wherein the fatty acid has an average of about 14 carbon atoms and the fatty alcohol has an average of about 16 carbon atoms per molecule.

8. The composition of Claim 6 wherein said (a) is
30 present in an amount of from about 0.15 to 0.35%, R^1 has an average of 18 carbon atoms, R^2 and R^3 are each alkyl groups having 1 carbon atom, R^4 is a benzyl group, and A is a chloride anion.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US90/00802

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) *

According to International Patent Classification (IPC) or to both National Classification and IPC

IPC(5): A61K 7/00, 7/06, 7/08

U.S. CL. 424/70

II. FIELDS SEARCHED

Minimum Documentation Searched †

Classification System

Classification Symbols

U.S.

514/938,943; 424/70; 252/357,547,DIG.13

Documentation Searched other than Minimum Documentation
to the Extent that such Documents are Included in the Fields Searched ‡

III. DOCUMENTS CONSIDERED TO BE RELEVANT †

Category *	Citation of Document, †† with indication, where appropriate, of the relevant passages ‡‡	Relevant to Claim No. ‡
X	US, A, 4,183,917 (IWAQ et al.) 15 January 1980 See Example III, Column 7 and Column 3, line 61.	1-8
Y	US, A, 4,654,207 (PRESTON) 31 March 1987 See the entire document.	1-8
A,P	US, A, 4,824,594 (HOEFFKES et al.) 25 April 1989 See the entire document.	1-8
A	US, A, 4,275,055 (NACHTIGAL et al.) 23 June 1981 See the entire document.	1-8

* Special categories of cited documents: ‡‡

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search

18 March 1990

Date of Mailing of this International Search Report

27 APR 1990

International Searching Authority

ISA/US

Signature of Authorized Officer

AMY HULINA

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.